



PRECLIS

WWW.RECIST.COM


PAREXEL[®]

YOUR JOURNEY. OUR MISSION.[™]



T 1.1

Welcome to the **RECIST 1.1** Quick Reference



*Eisenhauer, E. A., et al.
New response evaluation
criteria in solid
tumours: Revised
RECIST guideline
(version 1.1). Eur J
Cancer 2009;45:228-47.

Subject Eligibility

Only patients with measurable disease at baseline should be included in protocols where objective tumor response is the primary endpoint. Measurable disease is defined as the presence of at least one measurable lesion.

In studies where the primary endpoint is tumor progression (either time to progression or proportion with progression at a fixed date), the protocol must specify if entry is restricted to those with measurable disease or whether patients having non-measurable disease only are also eligible.

Methods of Assessment

The same method of assessment and the same technique should be used to characterize each identified and reported lesion at baseline and during follow-up.

- CT is the best currently available and reproducible method to measure lesions selected for response assessment. MRI is also acceptable in certain situations (e.g., for body scans but not for lung).
- Lesions on a chest X-ray may be considered measurable lesions if they are clearly defined and surrounded by aerated lung. However, CT is preferable.
- Clinical lesions will only be considered measurable when they are superficial and ≥ 10 mm in diameter as assessed using calipers. For the case of skin lesions, documentation by color photography, including a ruler to estimate the size of the lesion, is recommended.
- Ultrasound (US) should not be used to measure tumor lesions.

Methods of Assessment

- Tumor markers alone cannot be used to assess response. If markers are initially above the upper normal limit, they must normalize for a patient to be considered in complete response.
- Cytology and histology can be used in rare cases (e.g., for evaluation of residual masses to differentiate between Partial Response and Complete Response or evaluation of new or enlarging effusions to differentiate between Progressive Disease and Response/Stable Disease).
- Use of endoscopy and laparoscopy is not advised. However, they can be used to confirm complete pathological response.

Baseline Disease Assessment

All baseline evaluations should be performed as closely as possible to the beginning of treatment and never more than 4 weeks before the beginning of the treatment.

Measurable lesions

Must be accurately measured in at least one dimension (longest diameter in the plane of measurement is to be recorded) with a minimum size of:

- 10 mm by CT scan (CT scan slice thickness no greater than 5 mm; when CT scans have slice thickness >5 mm, the minimum size should be twice the slice thickness).
- 10 mm caliper measurement by clinical exam (lesions which cannot be accurately measured with calipers should be recorded as non-measurable).
- 20 mm by chest X-ray.

Baseline Disease Assessment

Measurable lesions

- **Malignant lymph nodes**

To be considered pathologically enlarged and measurable, a lymph node must be ≥ 15 mm in short axis when assessed by CT scan (CT scan slice thickness is recommended to be no greater than 5 mm). At baseline and in follow-up, only the short axis will be measured and followed.

- **Lytic bone lesions or mixed lytic-blastic lesions** with identifiable soft tissue components that can be evaluated by cross-sectional imaging techniques such as CT or MRI can be considered measurable if the soft tissue component meets the definition of measurability described above.
- **'Cystic lesions'** thought to represent cystic metastases can be considered measurable if they meet the definition of measurability described above. However, if non-cystic lesions are present in the same patient, these are preferred for selection as target lesions.

Baseline Disease Assessment

Non-measurable lesions

Non-measurable lesions are all other lesions, including small lesions (longest diameter <10 mm or pathological lymph nodes with 10 to <15 mm short axis), as well as truly non-measurable lesions. Lesions considered truly non-measurable include: leptomeningeal disease, ascites, pleural or pericardial effusion, inflammatory breast disease, lymphangitic involvement of skin or lung, abdominal masses/abdominal organomegaly identified by physical exam that is not measurable by reproducible imaging techniques.

- **Blastic bone lesions** are non-measurable.
- **Lesions with prior local treatment**, such as those situated in a previously irradiated area or in an area subjected to other loco-regional therapy, are usually not considered measurable unless there has been demonstrated progression in the lesion. Study protocols should detail the conditions under which such lesions would be considered measurable.

Target Lesions

- All measurable lesions up to a maximum of two lesions per organ and five lesions in total, representative of all involved organs, should be identified as **target lesions** and recorded and measured at baseline.
- Target lesions should be selected on the basis of their size (lesions with the longest diameter) and be representative of all involved organs, as well as their suitability for reproducible repeated measurements.
- All measurements should be recorded in metric notation using calipers if clinically assessed.

A sum of the diameters (longest for non-nodal lesions, short axis for nodal lesions) for **all target lesions** will be calculated and reported as the baseline sum diameters, which will be used as reference to further characterize any objective tumor regression in the measurable dimension of the disease. If lymph nodes are to be included in the sum, only the short axis will contribute.

Non-target Lesions

All lesions (or sites of disease) not identified as target lesions, including pathological lymph nodes and all non-measurable lesions, should be identified as **non-target lesions** and be recorded at baseline. Measurements of these lesions are not required and they should be followed as 'present', 'absent' or in rare cases, 'unequivocal progression'.

Response Criteria

Evaluation of target lesions

Complete Response (CR):

Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to <10 mm.

Partial Response (PR):

At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum of diameters.

Progressive Disease (PD):

At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this may include the baseline sum). The sum must also demonstrate an absolute increase of at least 5 mm.

Stable Disease (SD):

Neither sufficient shrinkage to qualify for PR nor sufficient increase to qualify for PD.

Response Criteria

Special notes on the assessment of target lesions

- **Lymph nodes** identified as target lesions should always have the actual short axis measurement recorded even if the nodes regress to below 10 mm on study. When lymph nodes are included as target lesions, the 'sum' of lesions may not be zero even if complete response criteria are met since a normal lymph node is defined as having a short axis of <10 mm.
- **Target lesions that become 'too small to measure'**. While on study, all lesions (nodal and non-nodal) recorded at baseline should have their actual measurements recorded at each subsequent evaluation, even when very small. However, sometimes lesions or lymph nodes become so faint on a CT scan that the radiologist may not feel comfortable assigning an exact measure and may report them as being 'too small to measure', in which case a default value of 5 mm should be assigned.

Response Criteria

Special notes on the assessment of target lesions

- **Lesions that split or coalesce on treatment.**
When non-nodal lesions 'fragment', the longest diameters of the fragmented portions should be added together to calculate the target lesion sum. Similarly, as lesions coalesce, a plane between them may be maintained that would aid in obtaining maximal diameter measurements of each individual lesion. If the lesions have truly coalesced such that they are no longer separable, the vector of the longest diameter in this instance should be the maximal longest diameter for the 'coalesced lesion'.

Response Criteria

Evaluation of non-target lesions

Complete Response (CR):

Disappearance of all non-target lesions and normalization of tumor marker levels. All lymph nodes must be non-pathological in size (<10 mm short axis).

Non-CR / Non-PD:

Persistence of one or more non-target lesion(s) and/or maintenance of tumor marker levels above normal limits.

Progressive Disease (PD):

Unequivocal progression of existing non-target lesions.

- **When patient has measurable disease.**
To achieve 'unequivocal progression' on the basis of the non-target disease, there must be an overall level of substantial worsening in non-target disease such that, even in presence of SD or PR in target disease, the overall tumor burden has increased sufficiently to merit discontinuation of therapy. A modest 'increase' in the size of one or more non-target lesions is usually not sufficient to qualify for unequivocal progression status.

Response Criteria

Progressive Disease (PD):

Unequivocal progression of existing non-target lesions.

- **When patient has only non-measurable disease.** There is no measurable disease assessment to factor into the interpretation of an increase in non-measurable disease burden. Because worsening in non-target disease cannot be easily quantified, a useful test that can be applied is to consider if the increase in overall disease burden based on change in non-measurable disease is comparable in magnitude to the increase that would be required to declare PD for measurable disease. Examples include an increase in a pleural effusion from 'trace' to 'large' or an increase in lymphangitic disease from localized to widespread.

Response Criteria

New lesions

The appearance of new malignant lesions denotes disease progression:

- The finding of a new lesion should be unequivocal (i.e., not attributable to differences in scanning technique, change in imaging modality or findings thought to represent something other than tumor, especially when the patient's baseline lesions show partial or complete response).
- If a new lesion is equivocal, for example because of its small size, continued therapy and follow-up evaluation will clarify if it represents truly new disease. If repeat scans confirm there is definitely a new lesion, then progression should be declared using the date of the initial scan.
- A lesion identified on a follow-up study in an anatomical location that was not scanned at baseline is considered a new lesion and disease progression.

Response Criteria

New lesions

It is sometimes reasonable to incorporate the use of FDG-PET scanning to complement CT in assessment of progression (particularly possible 'new' disease). New lesions on the basis of FDG-PET imaging can be identified according to the following algorithm:

Negative FDG-PET at baseline, with a positive FDG-PET at follow-up is PD based on a new lesion.

No FDG-PET at baseline and a positive FDG-PET at follow-up:

- If the positive FDG-PET at follow-up corresponds to a new site of disease confirmed by CT, this is PD.
- If the positive FDG-PET at follow-up is not confirmed as a new site of disease on CT, additional follow-up CT scans are needed to determine if there is truly progression occurring at that site (if so, the date of PD will be the date of the initial abnormal FDG-PET scan).
- If the positive FDG-PET at follow-up corresponds to a pre-existing site of disease on CT that is not progressing on the basis of the anatomic images, this is not PD.

Time Point Response

Table 1 provides a summary of the overall response status calculation at each time point for patients who have measurable disease at baseline.

Table 1. Time point response: Patients with target (+/- non-target) disease

Target lesions	Non-target lesions	New lesions	Overall response
CR	CR	No	CR
CR	Non-CR /non-PD	No	PR
CR	NE	No	PR
PR	Non-PD /or not all evaluated	No	PR
SD	Non-PD /or not all evaluated	No	SD
Not all evaluated	Non-PD	No	NE
PD	Any	Yes or No	PD
Any	PD	Yes or No	PD
Any	Any	Yes	PD

CR = Complete Response

PR = Partial Response

SD = Stable Disease

PD = Progressive Disease

NE = Inevaluable

Time Point Response

When patients have non-measurable (therefore non-target) disease only, Table 2 is to be used.

Table 2. Time point response: Patients with non-target disease

Non-target lesions	New lesions	Overall response
CR	No	CR
Non-CR/non-PD	No	Non-CR/non-PD ¹
Not all evaluated	No	NE
Unequivocal PD	Yes or No	PD
Any	Yes	PD

CR = Complete Response

PD = Progressive Disease

NE = Inevaluable

¹ Non-CR / non-PD is preferred over 'Stable Disease' for non-target disease since SD is increasingly used as an endpoint for assessment of efficacy in some trials. To assign this category when no lesions can be measured is not advised.

Confirmation

In non-randomized trials where response is the primary endpoint, confirmation of PR and CR is required to ensure responses identified are not the result of measurement error. This will also permit appropriate interpretation of results in the context of historical data where response has traditionally required confirmation in such trials.

However, in all other circumstances, (i.e., in randomized phase II or III trials or studies where stable disease or progression are the primary endpoints), confirmation of response is not required since it will not add value to the interpretation of trial results. However, elimination of the requirement for response confirmation may increase the importance of central review to protect against bias, in particular in studies which are not blinded.

In the case of SD, measurements must have met the SD criteria at least once after study entry at a minimum interval (in general not less than 6–8 weeks) that is defined in the study protocol.


Missing Assessments and Inevaluable Designation

When no imaging/measurement is done at all at a particular time point, the patient is not evaluable (NE) at that time point.

If only a subset of lesion measurements are made at an assessment, usually the case is also considered NE at that time point, unless a convincing argument can be made that the contribution of the individual missing lesion(s) would not change the assigned time point response. This would most likely happen in the case of PD.

RECIST 1.1

Frequently Asked Questions



*Compiled from RECIST 1.1 but inclusive of only those items that were not covered in the main body of the article

Q

When measuring the longest diameter of target lesions in response to treatment,

is the same axis that was used initially used subsequently, even if there is a shape change to the lesion that may have produced a new longest diameter?

1**A**

The longest diameter of the lesion should always be measured even if the actual axis is different from the one used to measure the lesion initially (or at a different time point during follow-up). The only exception to this is lymph nodes—per RECIST 1.1 the short axis should always be followed and as in the case of target lesions, the vector of the short axis may change on follow-up.

Q

Are RECIST criteria accepted by regulatory agencies?

2**A**

Many cooperative groups and members of the pharmaceutical industry were involved in preparing RECIST 1.0 and have adopted them. The FDA was consulted in their development and supports its use, though they don't require it. The European and Canadian regulatory authorities also participated and the RECIST criteria are now integrated in the European note for guidance for the development of anticancer agents. Many pharmaceutical companies are also using RECIST criteria. RECIST 1.1 was similarly widely distributed before publication.

Q

What if a single non-target lesion cannot be reviewed (for whatever reason)?

3

Does this negate the overall assessment?

A

Sometimes the major contribution of a single non-target lesion may be in the setting of CR having otherwise been achieved; failure to examine one non-target in that setting will leave you unable to claim CR. It is also possible that the non-target lesion has undergone such substantial progression that it would override the target disease and render the patient PD. However, this is very unlikely, especially if the rest of the measurable disease is stable or responding.

Q

A lesion which was solid at baseline has become necrotic in the center. How should this be measured?

4**A**

The longest diameter of the entire lesion should be followed. Eventually, necrotic lesions which are responding to treatment decrease in size. In reporting the results of trials, you may wish to report on this phenomenon if it is seen frequently since some agents (e.g., angiogenesis inhibitors) may produce this effect.

Q

If I am going to use MRI to follow disease, what is the minimum size for measurability?

5**A**

MRI may be substituted for contrast enhanced CT for some sites, but not lung.

The minimum size for measurability is the same as for CT (10 mm) as long as the scans are performed with a slice thickness of 5 mm and no gap. In the event the MRI is performed with thicker slices, the size of a measurable lesion at baseline should be two times the slice thickness. In the event there are inter-slice gaps, this also needs to be considered in determining the size of measurable lesions at baseline.

Q

Can PET-CT be used with RECIST?

6**A**

At present, the low dose or attenuation correction CT portion of a combined PET-CT is not always of optimal diagnostic CT quality for use with RECIST measurements. However, if your site has documented that the CT performed as part of a PET-CT is of the same diagnostic quality as a diagnostic CT (with IV and oral contrast) then the PET-CT can be used for RECIST measurements. Note, however, that the PET portion of the CT introduces additional data which may bias an investigator if it is not routinely or serially performed.

Q

A patient has a 32% decrease in sum cycle 2, a 28% decrease in cycle 4 and a 33% decrease in cycle 6.

7

Does confirmation of PR have to take place in sequential scans or is a case like this confirmed PR?

A

It is not infrequent that tumor shrinkage hovers around the 30% mark. In this case, most would consider PR to have been confirmed looking at this overall case. Had there been two or three non-PR observations between the two time point PR responses, the most conservative approach would be to consider this case SD.

To learn more about
how PAREXEL can
apply medical imaging
to your next oncology
program, contact us:

PAREXEL International

1 Federal Street

Billerica, MA 01821 USA

US: +1 978 313 3900

UK: +44 (0) 115 955 7333

Germany: +49 (0) 30 30 685 5075

E-mail: info@PAREXEL.com

www.recist.com

PAREXEL®
YOUR JOURNEY. OUR MISSION.™

© 2014 PAREXEL International Corporation.
All rights reserved.